

**Sternal - Gastric Aspecific Reflex (Str. G. A. R.) mean-intense digital pressure on any point of the spleen projection area– (sternal trigger points)**

Latency time (Lt) in seconds	Latency time after preconditioning (pause of 5 sec.)	MFR in seconds	fD & equilibria	EBD	Preconditioning	tCG	Diagnosis
Lt = 10	Lt = 20 [Lt = 20 if intense digital pressure on single sternal trigger point — absence of bone marrow disorder]	3 < MFR < 4 normal MFR, associated activation, outcome +	fD ≥ 3 (ideal value fD=3.81) stange attractor	Normal EBD physiological function	Type I Physiological tissue microvascular unit	<b>Negative tonic Gastric Contraction - tCG</b>	Health
Lt = 10	Lt < 20 [Lt = 0 if intense digital pressure on single sternal trigger point – presence of bone marrow disorder]	MFR = 4 compromised MFR, dissociated activation, outcome ±	2 < fD < 3 limit cycle	Normal, slightly modified EBD function, small number of pathological EBD	Type II A Intermediate tissue microvascular unit	<b>tonic Gastric Contraction - tGC</b> - local autoimmune syndrome - accompanied by gallbladder - and splenic contraction - decongestion: <b>positive tCG</b>	Myeloid Leukemia Inherited Real Risk
7 < Lt < 10	14 < Lt < 20 [Lt = 0 if intense digital pressure on single sternal trigger point – presence of bone marrow disorder]	4 < MFR ≤ 5 growing compromised MFR, dissociated activation, outcome ±	1 < fD ≤ 2 limit cycle	Modified EBD function, increasing number of pathological EBD	Type II B Intermediate tissue microvascular unit	<b>tonic Gastric Contraction - tGC</b> - local autoimmune syndrome - accompanied by gallbladder - and splenic contraction - decongestion: <b>positive tCG</b>	Myeloid Leukemia Inherited Real Risk in evolution
Lt ≤ 7	Lt < 14 [Lt = 0 if intense digital pressure on single sternal trigger point – presence of bone marrow disorder]	MFR > 5 absent MFR, dissociated activation, outcome –	fD = 1 fix point	Normal EBD function pathological, large number of pathological EBD	Type III Pathological tissue microvascular unit	<b>tonic Gastric Contraction - tGC</b> - local autoimmune syndrome - accompanied by gallbladder - and splenic contraction - decongestion: <b>positive tCG</b>	Overt Myeloid Leukemia

Table 1. Legend: MFR (Microcirculatory Functional Reserve); EBD (Endoarteriolar Blocking Device); CAD (Coronary Artery Disease; fD (fractal Dimension); Lt (Latency time)

PAY ATTENTION: In case of **lymphocytic leukemia**, the triggers point are those of spleen or lymph nodes, and the values of the above table are the same